

1. Mercier J.L., Dejean A. and Lenoir A, 1998. Limited Aggressiveness Among African Arboreal Ants sharing the same Territories: The Result of a Co- Evolutionary Process. *Sociobiology* 32(1).
2. Corbara B , 2000. Self organization in the division of labor in incipient societies of the Ant *Ectatomma rudium* (Formicidae: Ponerinae) : Emergence of an age polyethism, *Sociobiology* 36(3).
3. Yoshimoto T and Yamane S,1990. Food habits of some ant species inhabiting the Taisho lava of the Sakura –jima volcano, Japan, *Reports of the Faculty of Science Kagoshima University (Earth sciences and Biology)*, 23: 9-22.
4. Rajagopal D, Musthak Ali T.M. and Chakravarthy A.K, 1998. Ant species richness, across the altitudinal gradient in the western ghats of Karnataka, *ZOOS' PRINT*.
5. Veeresh G.K. and Musthak Ali T.M, 1990. The present status of our knowledge on Indian ants, *Social Insects*, 77-102.
6. Underwood. E .C , Quinn J.F, 2010. Response of ants and spiders to prescribed fire in oak woodlands of California. *J Insect Conserv* 14: 359-366.
7. Braschle B, Mahood K, Karenzi N, Gaston K.J, Chown S.L , 2009. Realizing a synergy between research and education: how participation in ant monitoring helps raise biodiversity awareness in a resource poor , *J Insect Conserv* 14: 19-30.
8. Castracani C, Grasso D. A, Fanfani A, Mori A, The ant fauna of Castelporziano presidential reserve ( Rome, Italy) as a model for the analysis of ant community structure in relation to environmental variation in Mediterranean ecosystems, *J Insect Conserv*, 14: 585.
9. Gotelli N.J, Ellison M, Dunn R.R and Sanders N.J, 2011. Counting ants: biodiversity sampling and statistical analysis for myrmecologists, *Myrmecol News* 15: 13-19
10. Peiris H.A.W.S., Dias R.K.S , 2010. Similarities and differences of ant communities in three types of habitats in the intermediate zone of Sri Lanka, *Proceedings of annual research symposium of University of Kelaniya*, 27pp.
11. Mooney K.A, Tillberg C.V , 2005. Temporal and Spatial variation to ant omnivory in pine forests, *Ecology*, 86(5): 1225-1235.
12. Levey D.J, Byrne M.M , Complex Ant – plant interactions: rain forest ants as Secondary dispersers and post- dispersal seed predators, *Ecology*, 74:1802-1812.
13. Basset Y,2001. Invertebrates in the canopy of tropical rain forests How much do we really know? *Plant Ecology* 153: 87-107.
14. Basset Y, Aberlenc H.P and Delvare G,1992. Abundance and stratification of foliage arthropods in a lowland rain forest of Cameroon, *Ecological Entomology*, 17: 310-318.
15. Veeresh G.K, Musthak Ali T.M and Kumar A.R.V, Ants as balancers in nature pest problems, *Biological control of social forest and plantation crops Insects*, 122-135.
16. Basset Y, Samuelson V, Allison A and Miller S.E, 1996. How many species of host specific insects feed on a species of tropical tree? , *Biological Journal of the Linnean society*, 59: 201-216.
17. Baur A, Sanetra M, Chalwatizis N, Buschinger A and Zimmermann F.K,1996. Sequence comparisons of the international transcribed spacer region of ribosomal genes support close relationships between parasitic ants and their respective host species, *Ins. Soc.* 43: 53-67.
18. Yamane S, Harada Y, Yano M, 1985. Ant fauna of Tanega- shima island, the northern Ryukyus, *Mem. Kagoshima Univ. Res. Center s. Pac*, 6(1):166-173.

19. YAMANE S, IWAI T, WATANABE .H, 1994. Ant fauna of the Tolara islands, Northern Ryukyus, Japan, WWF Japan Science Report 2(2): 311-327.
20. IVANOV K, MILLIGAN J, KEIPER J, 2010. Efficiency of the Winkler method for extracting ants from temperate- forest litter, *Myrmecological News* 13:73-79.
21. Lachaud J.P and Lachaud G.P, 2012. Diversity of species and behaviour of Hymenopteran Parasitoids of Ants, A review. *Psyche*, 2012:24.
22. Boulton M, Davies K.F and Ward P.S, 2005. Species richness, abundance, and composition of Ground- dwelling ants in Northern California grasslands: Role of plants, soil, and grazing, April, *Environmental entomology*, 34(1):96-104.
23. Orivel J, Dejean A, Errard C, 1997. Ant gardens: interspecific recognition in parabiocotic ant species, *Behav. Ecol sociobiol* 40: 87-93.
24. Mercier J.L, Lenoir A, 1999. A new case of jumping behavior in ants, as part of the foraging strategy, *C.R. Acad. Sci. Paris, sciences de la vie/ Life sciences* 322: 661-667.
25. Mercier J.L and Dejean A, 1996. Ritualized behavior during competition for food between two Formicinae., *Ins. Soc.* 43: 17-29.
26. Vienne C, Errard C and Lenoir A (1997), Influence of the queen on worker behavior and queen recognition behavior in ants. *Ethology* 104:431-446.
27. Lenoir A, Fresneau D, Errard C and Hefetz A, 1999. Individuality and colonial identity in ants: the emergence of the social representation concept, *Information processing in social insects*, Birkhäuser Basel 3 :219-237.
28. Keller R.A, 2000. Cladistics of the tribe *Ectatommini* (Hymenoptera: Formicidae): A reappraisal, *Insects systematic and evolution* 31:4, 60-70.
29. Torres J.A, Snelling R.R and Canals M, 2001. Seasonal and Nocturnal periodicities in Ant Nuptial flights in the tropics ( Hymenoptera: Formicidae), *Sociobiology J Insect Conserv*, 37(3B): 601-626.
30. Widodo E.S, Naito T, Mohamed M, and Hashimoto Y, 2004. Effects of selective logging on the arboreal ants of a Bornean rainforest. *Entomological Science* 7: 341-349.
31. Widodo E.S, Mohamed M and Hashimoto Y, 2001. Canopy ant diversity assessment in the fragmented rainforest of Sabah, East Malaysia., *Nature and human activities* 6:13-23.
32. Lubertazzi D and Tschinkel W.L, 2003. Ant community change across a ground vegetation gradient in north Florida's longleaf pine flatwoods. *Journal of Insect Science*, 3(21):17.
33. Mackay W.P, Castro C.L and Fernandez F, 2002. A new, high altitude Colombian Species of the ant genus *Camponotus* with *Dimorphic* males and females (Hymenoptera: Formicidae). *Sociobiology*, 40:2.
34. Mackay W.P and Mackay E.E, 2000. A note in a response to Mr. Roy Snelling's Review of the *Camponotus montivagus* Complex ( Hymenoptera: Formicidae). *Sociobiology*, 36:3.
35. Mackay W.P, 1998. A remarkable new species of *Paratrechina* (Hymenoptera: Formicidae) from the state of Colima, Mexico. *Pan-Pacific Entomologist*, 74(2): 99-101.
36. Watanabe H, Yamane S, 1999. New species and new status in the genus *Aphaenogaster* ( formicidae) from Japan, Yamane, S., S. Ikudome and Terayama M. Identification guide to the Aculeata of the Nansei Islands, Japan, 728-736 pp.
37. Mackay W.P and Buschinger A, 2002. A new species of the Ant Genus *Leptothorax* ( Subgenus *Myrafant*) from Alberta, Canada, *Sociobiology* 40:3.

38. Stiles J.H and H.Jones R.H, 1998.Distribution of the red imported fire ant, *Solenopsis invicta*, in road and powerline habitats. *Landscape Ecology*, 335: 335-346.
39. Waser N.M, 1998.Task- Matching and Short- Term size shifts in foragers of the Harvester Ant, *Messor pergandei* ( Hymenoptera: Formicidae). *Journal of Insect behavior*, 11:3.
40. Babu M. J, Ankolekar S.M and Rajashekhar K.P, 2011.Castes of the weaver ant *Oecophylla smaragdina* ( Fabricius) differ in the organization of sensilla on their antennae and mouthparts. *Current Science*,101:6.
41. Ward P.S, Brady S.G, Fisher B.L and Sahultz T.R, 2015.The evolution of myrmicine ants: phylogeny and biogeography of a hyperdiverse ant clade( Hymenoptera: Formicidae), *systematic Entomology*, 40:61-81.
42. Marcela P, Abu Hassan A., Nurita A.T. and Thevan K,2012.Colony structure of the Weaver Ant, *Oecophylla smaragdina* ( Fabricius) ( Hymenoptera: Formicidae). *Sociobiology*,59:1.
43. Wood W.F, Palmer T.M, Staton M.L.(2006), Volatiles in the mandibular gland of tetraponera penzigi: A plant ant of the whistling thorn acacia. *Biochemical systematics and ecology*, 34:536-538.
44. Jacquemin J, Sonet G, Bourguignon T, Evans T A, Delsinne T.(2015), Second record and DNA barcode of the ant *Tyrannomyrmex rex* Fernandez ( Hymenoptera: Formicidae: Myrmicinae). *Sociobiology* 62(2): 276-280.
45. Braekman J.C, Daloze D, Pasteels J.M, Hecke P.V, 1987.Tetraponerine-8, an Alkaloidal contact poison in a Neoguinean Pseudomyrmecine ant, *Tetraponera* sp. *Z.Naturforsch.* 42c:627-630.
46. Weber N.A, 1949.The functional significance of dimorphism in the African ant, *Oecophylla*., *Ecology*,30:3.
47. Kohout R.J, 1998.New synonyms and nomenclatural changes in the ant genus *Polyrhachis* Fr. Smith( Hymenoptera: Formicidae: Formicinae). *Memoirs of the Queensland Museum* 42(2): 505-531.
48. Ogata K and Onoyama K, 1998. A revision of the Ant Genus *Smithistruma* brown of Japan, with description of four new species ( Hymenoptera: Formicidae), *Entomological Science* 1(2) : 277-287.
49. Terayama M and Yamane S, 2000. *Lasiomyrma*, a new stenammine Ant Genus from Southeast Asia ( Hymenoptera: Formicidae), *Entomological Science*, 3(3): 523-527.
50. Samarawickrama V.A.M.P.K, Janananda M.D.B.G, Ranawana K.B., and Smith A, 2005.Study of the distribution of the Genus *Poecilotheria* of the family Theraphosidae in Sri Lanka, *Cey. J. Sci. ( Bio. Sci.)* 34: 75-86.
51. Shamsadini S, 2003. Localized scalp hair shedding caused by *Pheidole* ants and overview of similar case reports, *Dermatology online Journal*, 9:3.
52. Eguchi K, 2000.Two new *Pheidole* species with a 5- segmented antennal club ( Hymenoptera: Formicidae) , *Entomological Science*3(4): 687-692.
53. Nowbahari E et al, 2000. Polymorphism and Polyethism in the Formicinae Ant *Cataglyphis niger* ( Hymenoptera) , *Sociobiology* 36:3.
54. Camargo R X, Oliveira P S.2012.Natural history of the Neotropical arboreal ant, *Odontomachus hastatus*: Nest sites, foraging schedule, and diet, *journal of insect science*,12( 48).

55. Bolton B, 1991. New myrmicine ant genera from the oriental region (Hymenoptera: Formicidae), *Systematic Entomology* 16, 1-13.
56. Mercier J.I and Lenoir A, 1999. Individual flexibility and choice of foraging strategy in *Polyrhachis laboriosa* F. Smith (Hymenoptera, Formicidae), *Insectes soc*, 46: 267-272.
57. Terayama M, Ito F and Gobin B, 1998. Three new species of the Genus *Acanthomyrmex* Emery (Hymenoptera: Formicidae) from Indonesia, with Notes on the reproductive Caste and Colony Composition, *Entomological Science* 1(2): 257-264.
58. Maeyama T, Machida M and Terayama M, 1999. The ant-parasitic genus *Rhipipalloidea giraulti* (Hymenoptera: Eucharitidae), with a description of a new species, *Australian Journal of Entomology*, 38: 305-309.
59. Mercier J.L, Lenoir A, Dejean A, 1997. Ritualised versus aggressive behaviours displayed by *Polyrhachis laboriosa* (F. Smith) during intraspecific competition, *Behavioural processes*, 41: 39-50.
60. Sanetra M, Gusten R and Schulz A, On the taxonomy and distribution of Italian *Tetramorium* species and their social parasites (Hymenoptera Formicidae). *Mem. Soc. Entomol. Ital*, 77: 317-357.
61. Villet M.H, 1999. Reproductive behavior of *Plectroctena mandibularis* F. Smith (Hymenoptera: Formicidae), a ponerine ant with ergatoid queens, *African Entomology*, 7: 2.
62. Dubois M.K, 2000. *Monomorium minimum* species group: Gyne number and longevity (Hymenoptera: Formicidae), *Entomological news*, 111(1): 13-20.
63. Onoyama K, 1998. Taxonomic notes on the ant Genus *Crematogaster* in Japan (Hymenoptera: Formicidae), *Keiichi, Entomological science*, 1(2): 227-232.
64. Eguchi K, 2001. Sorting Asian *Pheidole* at Species level, and establishing colony-based reference collections of Ants (Hymenoptera: Formicidae), *ANeT Domestic workshop in Thailand*.
65. Gyung K.K and Eguchi K, 2001. Taxonomy of "*Deromyrma*" of the genus *Aphaenogaster* in the Indo-Chinese and Indo-Malay Subregions (Hymenoptera: Formicidae, Myrmicinae): A prompt result, *Proceedings of ANeT in Hanoi*.
66. Mackay W.P and Mackay E.E, 1997. A revision of the Neotropical Ants of the Montivagus species complex, Genus *Camponotus*, Subgenus *Myrmentoma* (Hymenoptera: Formicidae), *Sociobiology*, 30(3).
67. Gaglio M D, Mackay W P, Osorio E A, and Iniguez I, 1998. Nest population of *Pogonomyrmex salinus* Harvester Ants (Hymenoptera: Formicidae), *Sociobiology*, 32: 3.
68. Mackay W P, A new army ant from Southern Texas (Hymenoptera: Formicidae), *The Southwestern naturalist*, 43(3): 333-335.
69. Fernando Fernandez C. and Mackay W P, 2003. The Myrmicine ants of the *Adelomyrmex laevigatus* species complex (Hymenoptera: Formicidae), *Sociobiology*, 41: 3.
70. Sotrz SR, Tschinkel WR., 2004. Distribution, spread, and ecological associations of the introduced ant *Pheidole obscurithorax* in the Southeastern, United States, *Journal of insect science*, 4: 12.
71. Ogata K, Terayama M and Masuko K, 1995. The ant genus *Leptanilla*: discovery of the worker-associated male of *L. japonica*, and a description of a new species from Taiwan (Hymenoptera: Formicidae, Leptanillinae), *Systematic Entomology*, 20: 27-34.

72. Eguchi K, 2001. A revision of the Bornean Species of the Ant Genus *Pheidole* ( Insecta: Hymenoptera: Formicidae, Myrmicinae), *Tropics*, 2: 1-154.
73. Eguchi K, Bui T V & Yamane S, 2004. A preliminary study on foraging distance and nesting sites of ants in Indo-Chinese lowland vegetation ( Insecta, Hymenoptera: Formicidae). *Sociobiology*, 43:3.
74. Fuminori ITO, Yamane S, Eguchi K, Noerdjito W A, Kahono S, Tsuji K, Ohkawara K, Yamauchi K, Nishida T, Nakamura K, 2001. Ant species diversity in the Bogor botanic garden , West Java, Indonesia, with descriptions of two new species of the Genus *Leptanilla* (Hymenoptera: Formicidae), *TROPICS* , 10(3): 379-404.
75. Tsuji K, Egashira K and Holldobler B, 1999. Regulation of worker reproduction by direct physical contact in the ant *Diacamma* sp. From Japan. *Animal Behaviour*, 58:337-343.
76. Yoshimura M and B L Fisher, 2012. A revision of male ants of the Malagasy region (Hymenoptera: Formicidae): Key to subfamilies and treatment of the genera of Ponerinae, *PLoS ONE* 7(3): e33325.
77. Wetterer J K, 2015. Geographic origin and spread of cosmopolitan ants (Hymenoptera: Formicidae), *HALTERES*, 6: 66-78.
78. McGlynn T .P, 1999. The worldwide transfer of Ants: Geographical distribution and Ecological Invasions, *Journal of biogeography*, 26(3):535-548.
79. Holway D.A, Lach L, Suarez A.V, Tsutsui N D, and Case T.J, 2002. The causes and consequences of Ant invasions, *Annual reviews*, 33:181-233.
80. Gunawardene N R, Majer J D & Edirisinghe J P, Diversity and richness of ant species in a lowland wet forest reserve in Sri lanka, *ASIAN MYRMECOLOGY*, 2: 71-83.
81. Brady S.G, Fisher B.L, Schultz T.R and Ward P.S, 2014. The rise of army ants and their relatives: diversification of specialized predatory doryline ants, *BMC Evolutionary Biology*:1471-2148.
82. Ward P.S, Blaimer B.B and Fisher B.L, 2016. A revised phylogenetic classification of the ant subfamily Formicinae (Hymenoptera: Formicidae), with resurrection of the genera *Colobopsis* and *Dinomyrmex*, *ZOOTAXA*, 343- 357.
83. Ward P.S, 2007. Phylogeny, classification, and species-level taxonomy of ants (Hymenoptera: Formicidae), *ZOOTAXA*, 549-563.
84. Mezger D and Moreau C.S, 2015. Out of South-East Asia: phylogeny and biogeography of spiny ant genus *Polyrhachis* Smith (Hymenoptera: Formicidae), *Systematic Entomology*, 41(2): 369-378.
85. Lapolla J.S, 2009. Taxonomic revision of the Southeast Asian Ant Genus *Euprenolepis*, *ZOOTAXA*, 2046: 1-25.
86. Dias R.K.S and Kosgamage K.R.K. A., 2012. Occurrence and species diversity of ground dwelling worker ants (Family : Formicidae) in selected lands in the dry zone of Sri Lanka, *J. Sci. Univ. Kelaniya*:55-72.
87. Conceicao E.S.D, Jacques H.C. Delabie E Antonio De O. Costa Neto, 2004. The Entomophily of the Coconut tree in question: The evaluation of pollen transportation by Ants (Hymenoptera: Formicidae) and bees (Hymenoptera: apoidea) in florescence, *Neotropical Entomology*, 33(6): 679-683.

88. Conceicao E.S.D, Moura J.I.L, costa-Neto A. D. O. and Delabie J.H.C, 2014. Choice of food resource by Bees and Ants (Hymenoptera: Apoidea, Formicidae) on Coconut tree inflorescences (*Cocos nucifera*, Arecaceae), *Annals de la Societe entomologique de France (N.S): International Journal of Entomology*, 50(1): 6-12.
89. Conceicao E.S.D, Delabie J.H.C, Lucia T.M.C.D, Costa-Neto A.D.O and Majer J.D, 2014. Structural changes in arboreal assemblages (Hymenoptera: Formicidae) in an age sequence of Cocoa plantations in the South-east of Bahia, Brazil , *Austral entomology*, 54 (3):315-324.
90. Folgarait P.J.F, 1998. Ant biodiversity and its relationship to ecosystem functioning: a review, *Biodiversity and conservation*, 7:1221-1244.
91. Jacquemin J, Drouet T, Delsinne T, Roisin Y, Leponce M, 2012. Soil properties only weakly affect subterranean ant distribution at small spatial scales, *Soil properties only weakly affect subterranean ant distribution at small spatial scales*, *Applied soil ecology*, 62: 163-169.
92. Geralch J, Samways M, Pryke J, 2013. Terrestrial invertebrates as bioindicators: an overview of available taxonomic groups, *Journal of Insect Conservation* ,17: 831-850.
93. Mackay W.P, 1998. Dos especies nuevas de hormigas de la tribu Attini de Costa Rica y Mexico: *Mycetosoritis vinsoni* y *Mycocepurus curvispinosus* (Hymenoptera: Formicidae), *Rev. Biol. Trop*, 46(2): 421-426.
94. Forel A, 1912. Descriptions provisoires de genres, sous-genres et especes de Formicides des Indes orientales, *REVUE SUISSE DE ZOOLOGIE*, 20(15): 762-774.
95. Pacheco R, Vasconcelos H.L, Groc S, Gabriela P. Camacho G.P, Frizzo T.L.M, 2013. The importance of remnants of natural vegetation for maintaining ant diversity in Brazilian agricultural landscapes, *Biodiversity Conservation*, 22:983.
96. Peiris H.A.W.S. and Dias R.K.S, Similarities and differences of ant communities in three types of habitats in the intermediate zone of Sri Lanka, *Proceedings of Annual research symposium of University of Kelaniya*, 127 pp.
97. Ganehiarachchi G.A.S.M. and Fernando I.V.S, 2000. *Parachrysocharis javensis* Girault (Hymenoptera: Eulopidae)- An effective egg parasitoid of Sugarcane planthopper *Pyrilla perpusilla* in the wet zone of Sri Lanka, *Vidyodaya journal of Science*, 9:129-135.
98. Ganehiarachchi G.A.S.M. and Fernando I.V.S, 2000. Natural enemies of Sugarcane planthopper *Pyrilla perpusilla* walker (Homoptera: Lophopidae) in the wet zone of Sri Lanka, *Journal of national Science foundation Sri Lanka*, 28(3):205-213.
99. Mackay W.P and Mackay E.E, 1997. A revision of the Neotropical ants of the montivagus species complex, Genus *Camponotus*, subgenus *Myrmentoma* (Hymenoptera: Formicidae), *Sociobiology* 30(3 ): 319-334.
100. Mackay W.P, 1997. A revision of the Neotropical ants of the Genus *Camponotus*, Subgenus *Myrmostenus* (Hymenoptera: Formicidae), *PROC. ENTOMOL. SOC. WASH.* , 99(1):194-203.
101. Mackay W.P and Mackay E. E, 1997. Una Nueva especie de Hormiga del Genero *Trachymyrmex* (Hymenoptera: Formicidae) del Estado de Chihuahua, Mexico, *Sociobiology*, 30(1).
102. Kim J.K and YAMANE S, 2001. A Revision of *Eumenes* Latreille (Hymenoptera: Vespidae) from the far East Asia, with descriptions of one new species and one new subspecies, *Entomological Science*, 4(2): 139-155.

103. Mackay W.P, Mena R, Gardea J, and Pingatore N, 1997.Lack of Bioaccumulation of heavy metals in an arthropod community in the Northern Chihuahuan desert, JOURNAL OF THE KANSAS ENTOMOLOGICAL SOCIETY, 70(4): 329-334.
104. Mackay W.P, Mena R, Pingatore J R N E, Redetzke K, Freeman C.E, Newman H, Gardea J, and Navarro H, 1998. Seasonal changes in concentration and distribution of heavy metals in Creosotebush, *Larrea Tridentata* (Zygophyllaceae), Tissues in the El Paso, TX/Ciudad Juarez, Mexico area, SIDA 18(1):287-296.
105. Gyung K and EGUCHI K, 2001. Taxonomy of “*Deromyrma*” of the genus *Aphaenogaster* in the Indo- Chinese and Indo-Malay subregions (Hymenoptera, Formicidae, Myrmicinae) : a prompt result, Proceedings of ANet.
106. Chao A, Chazdon R.L, Colwell R.K and Shen T.J, 2005.A new statistical approach for assessing similarity of species composition with incidence and abundance data, Ecology letters,8:148-159.
107. Ward P.S, Brady S.G, Brian L. Fisher B.L and Schultz T.R, 2015.The evolution of myrmicine ants: phylogeny and biogeography of a hyperdiverse ant clade (Hymenoptera: Formicidae) , Systematic Entomology, 40:61-81.
108. Lapolla J.S, Brady S.G and Shattuck S.O, 2010. Phylogeny and Taxonomy of the *Prenolepis* genus-group of Ants (Hymenoptera: Formicidae), Systematic Entomology, 35:118-131.
109. KRONAUER D.J.C, 2009.Recent advances in army ant biology (Hymenoptera : Formicidae) , Myrmecological news, 12: 51-65.
110. Saranat E.M, Fischer G, Guenard B, Economo E.P, 2015.Introduced *Pheidole* of the world: taxonomy, biology and distribution, Zookeys, 543: 1-109.
111. SCHMIDT C.A. & SHATTUCK S.O, 2014. The higher classification of the ant subfamily Ponerinae (Hymenoptera : Formicidae), with a review of Ponerine Ecology and behavior , Zootaxa 3817 (1) : 001-242.
112. Fischer G, Azorsa F, Fisher B.L,2014.  
The ant genus *Carebara* westwood (Hymenoptera, Formicidae) : synonymisation of Pheidologeton Mayr under *Carebara*, establishment and revision of the *C. polita* species group, Zookeys 438: 57-112.
113. Ward P.S, 2001.Change in marine communities: An approach to statistical Analysis and interpretation, Taxonomy, phylogeny and biogeography of the ant genus *Tetraponera* (Hymenoptera : formicidae) in the oriental and Australian regions, Invertebrate Taxonomy,15: 589- 665.
114. Garcia F.H, Fisher B.L, 2015. Taxonomy of the hyper-diverse ant genus Tetramorium Mayr in the Malagasy region (Hymenoptera, Formicidae, Myrmicinae) – first record of the *T. setigerum* species group and additions to the Malagasy species groups with an updated illustrated identification key, Zookeys ,512 : 121-153.
115. EGUCHI K, 2001. A revision of the Bornean species of the Ant genus *Pheidole* (Insecta : Hymenoptera : Formicidae : Myrmicinae) , TROPICS, 2:1-154.
116. Bolton B , 1997. Identification guide to the ant genera of the world, Cambridge university press, pp 232.
117. Shattuck S.O, 1992. Higher classification of the ant subfamilies Aneuretinae, Dolichoderinae and Formicinae (Hymenoptera : Formicidae), Systematic entomology, 17:199-206.

118. Karunarathna D.A.G.N.B. and Karunaratne W.A.I.P, 1893. Two new localities of Sri Lankan relict ant *Aneuretus simony emery* , (Formicidae : Aneuretinae) with the very first record in the intermediate zone, Journal of threatened Taxa, 5 (11) : 4604-4607.
119. Wilson E.O, Eisner T, Wheeler G.C. and Wheeler J, 1956. *Aneuretus simony emery* , A major link in Ant evolution, Bulletin of the museum of comparative Zoology, 115:3.
120. Jayasuriya A.K. and Traniello J.F.A., The biology of the primitive Ant *Aneuretus simony (emery)* (Formicidae : Aneuretinae) distribution, abundance, colony structure, and Foraging ecology, Insectes Sociaux, 32(4 ):363-374.
121. Traniello J.F.A, Jayasuriya A.K , 1985. The biology of the primitive Ant *Aneuretus simony (emery)* (Formicidae : Aneuretinae) II The social ethogram and division of labor , Insectes sociaux , 32(4):375-388.
122. Andersen A N , 2002. Common names for Australian ants (Hymenoptera : formicidae) , Australian journal of Entomology 41:285-293.
123. Ranil R.H.G. and Pushpakumara D.M.N.G, Occurrence of *Drynaria sparsisora* (Desv.) T. Moore, in the lower Hantana area, Sri Lanka, J.Natn.Sci.Foundation Sri Lanka 36(4), 331-334.
124. Currie C.R, Poulsen M, Mendenhall J, Boomsma J.J, Billen J, Coevolved Crypts and Exocrine glands support mutualistic Bacteria in Fungus-growing ants, Science reprint, 311:81-83.
125. Billen J, 2009. Occurrence and structural organization of the exocrine glands in the legs of ants, Arthropod structure and Development, 38:2-15.
126. Billen J, A survey of the glandular system of fire ants, Applied Myrmecology, 8:86-94.
127. Billen J, 2011.Exocrine glands and their key function in the communication system of social Insects, Formosan Entomol, 31: 75-84.
128. Billen J,2006.Signal variety and communication in social insects, PROC.NETH.ENTOMOL.SOC.MEET, 17:9-25.
129. Annette Van Oystaeyen A.V, Ricardo Caliari Oliveria R.C, Luke HolmanJelle S. Van Zweden L.H.S.V, Carmen Romero C, Cintia A. Oi C.A, Patrizia D’Ettorre P.D, Mohammadreza Khalesi M, Johan Billen J, Felix Wackers F, Jocelyn G. Millar J.G, Tom Wenseleers T, 2014. Conserved class of queen pheromones stops Social insect workers from reproducing, SCIENCE VOL 343:287-290,
130. YAMANE S, 1999. A new species of the genus *Pristomyrmex* MAYR from Japan, and a proposal of a New Synonym of species in the Genus *Camponotus* MAYR (Hymenoptera ; Formicidae), Mem. Myrmecology. Soc. Jpn., 1:17-24,
131. Ng’endo R.N, Osiemo Z.B, Brandl R,2012. DNA barcodes for species identification in the hyperdiverse ant genus *Pheidole* (Formicidae: Myrmicinae), Journal of Insect Science13(27) :1536-2442.
132. Lapolla J.S, Brady S.G and Shattuck S.O, 2010. Phylogeny and taxonomy of the *Prenolepis* genus-group of ants (Hymenoptera: Formicidae), Systematic entomology, 35: 118-131.
133. Blaimer B.B, 2012. A subgeneric revision of *Crematogaster* and discussion of regional species-groups (Hymenoptera: Formicidae) , Zootaxa 3482: 47-67.
134. Amarasinghe H.E, 2010. Species composition and nesting habits of ants in a hill-country home garden in Sri Lanka, ASIAN MYRMECOLOGY, 3:9-20.
135. Bharti H, Guenard B, Bharti M, Economo E.P, 2016. An updated checklist of the ants of India with their specific distributions in Indian states (Hymenoptera, Formicidae), Zookeys, 551: 1-83.



136. Ward P. S, Brady S.G, Fisher B. L. and Schultz T. R, 2015. The evolution of myrmicine ants : phylogeny and biogeography of a hyperdiverse ant clade (Hymenoptera: Formicidae), *Systematic Entomology* 40:61-81.
137. Wheeler G.C. and Wheeler J, 1976. Ant larvae: Review and Synthesis, *Memories of the Entomological Society of Washington*, 07: 1-108.
138. Wappler T, Dlussky G.M, Reuter M, 2009. The first fossil record of *Polyrhachis* (Hymenoptera: Formicidae: Formicinae) from the upper Miocene of Crete (Greece) *PALAONTOLOGISCHE ZEITSCHRIFT* ,83: 431-438.
139. Nest structure variation and morphology of adult castes of weaver ants *Polyrhachis* (Crytomyrma) *rastellata* (Latreille, 1802) (Hymenoptera: Formicidae) in Sri Lanka, Dias R.K.S, 2015. *Caucasian Entomological Bull.* 11(1): 125-130.
140. Dorow W.H.O, 1995. Revision of the ant genus *Polyrhachis* Smith, 1857 (Hymenoptera: Formicidae) on subgenus level with keys, checklist of species and bibliography, *Courier Forsch-Inst. Senckenberg*, 185: 1-113.
141. Hung A.C.F, A revision of the Ant genus *Polyrhachis* at the subgeneric level (Hymenoptera: Formicidae) , *Trans. Amer. Ent. Soc.*,93:395-422.
142. Dorow W.H.O, Maschwitz U and Rapp S, 1990. The natural history of *Polyrhachis* (Myrmhoptla) *muelleri* Forel 1893 (Formicidae Formicinae), a weaver ant with mimetic larvae and an unusual nesting behavior, *Tropical Zoology*, 3: 181-190.
143. Dorow W.H.O. and Kohout R.J, 1995. A review of the subgenus *Hemioptica* Roger of the genus *Polyrhachis* Fr. Smith with description of a new species (Hymenoptera: Formicidae: Formicinae), *Zool.Med.Leiden* ,69:94-104.
144. Robson S.K.A and Kohout R.J, 2005. Evolution of nest-weaving behavior in arboreal nesting ants of the genus *Polyrhachis* Fr.smith (Hymenoptera: Formicidae), *Australian Journal of Entomology*, 44:164-169.
145. Dorow W.H.O. and Maschwitz U, Robson S.K.A and Kohout R.J, 2005. The Arachne group of *Polyrhachis* (Formicidae, Formicinae): Weaver ants Cultivating Homoptera on Bamboo, *Insectes Sociaux*, 37(1): 73-89.
146. Wheeler G.C and Wheeler J, 1990. Larvae of the Formicinae ant genus *Polyrhachis*, *American Entomological Society*, 116 (3): 753-767.
147. C. Liefke C, Dorow W.H.O, Holldobler B and Maschwitz U, 1998. Nesting and food resources of syntopic species of the ant genus *Polyrhachis* (Hymenoptera, Formicidae) in west-Malaysia, *Insectes soc:* 45, 411-425.
148. Kohout R.J and Taylor R.W, 1990. Notes on Australian ants of the genus *Polyrhachis* FR.Smith, with a synonymic list of the species (Hymenoptera: Formicidae: Formicinae) , *Memories of the Queensland museum*, 28(2): 509-522.
149. Chen ZL, Zhou SY, YE DD, Chen Y, Lu CW, 2013. Molecular phylogeny of the ant subfamily Formicinae (Hymenoptera, Formicidae) from China based on Mitochondrial genes, *Sociobiology* 60(2): 135-144.
150. Pfeiffer M, Mezger D & Dyckmans J, 2014. Trophic ecology of tropical leaf litter ants (Hymenoptera: Formicidae) – a stable isotope study in four types of Bornean rain forest, *Myrmecological news*, 19:31-41.

151. Peeters C, 2014. Convergent evolution of wingless reproductives across all subfamilies of ants, and sporadic loss of winged queens (Hymenoptera: Formicidae), *Myrmecological news*, 19:31-41.
152. Shattuck S.O and Barnett N.J, 2006. Australian species of the ant genus *Diacamma* (Hymenoptera : Formicidae), *Myrmecologische Nachrichten*, 8:13-19.
153. Zweden J.S.V, Carew M.E., Henshaw M.T, Robson S.K.A. and Crozier R.H, 2007. Social and genetic structure of a supercolonial weaver ant, *Polyrhachis robsoni* , with dimorphic queens, *Insect. Soc.* 54:34-41.
154. New species of the *Polyrhachis* (*Myrma*) *parabiotica* species group (Hymenoptera : Formicidae: Formicinae) from the Philippines, Kohout R.J, 2006. *Australian Entomologist*, 33(3): 155-163.
155. Schoeters E and Billen J, 1993. Anatomy and fine structure of the Metapleural gland in *Atta* (Hymenoptera, Formicidae), *Belg. J. Zool*,123:67-35.
156. Nascimento R.R.D., Schoeters E, Morgan E.D, Billen J, and Stradling D.J, 1996. Chemistry of Metapleural Gland Secretions of Three Attine ants, *Atta sexdens rubropilosa*, *Atta cephalotes*, and *Acromyrmex octospinosus* (*Hymenoptera:Formicidae*) , *Journal of Chemical Ecology*, 5:987-1000.
157. BILLEN\* J.P.J and VAN BOVEN J.K.A, The metapleural gland in old world army ants: a morphological and ultrastructural description (hymenoptera, Formicidae), *Revue de Zoologie africaine*, 101:32-41.
158. Schoeters E and Billen J, 1992. Morphological and ultrastructural study of the metapleural gland in *Dicamma* (Hymenoptera, Formicidae), *Biology and Evolution of Social Insects* (J. Billen, Ed), 239-247.
159. Chemical composition and Function of Metapleural gland secretion of the Ant, *Crematogaster deformis* Smith (Hymenoptera: Myrmicinae), Attygalle A.B., Siegel B, Vostrowsky O, Bestmann H.J, and Maschwitz U, 1989. *Journal of Chemical Ecology*,15(1):317-328.
160. Fitzgerald KT, Flood AA, 2006. Hymenoptera Stings, *Clin Tech small Anim Pract*, 21(4): 194-204.
161. Billen J, 2009. Occurance and structural organization of the exocrine glands in the legs of ants, *Arthropod structure and Development*,38: 2-15.
162. Billen J, 2008. A novel exocrine gland in the trochanter of ant legs, *Acta Zoologica* (Stockholm), 89: 201-204.
163. MARTIN S.J, DILS V, BILLEN J, 2005. Morphology of the Dufour gland within the honey bee sting gland complex, *APIDOLOGIE* 36:543-546.
164. Morgan E.D, Jungnickel H, Billen J, Ito F, Bergman J, Gobin B, 2008. Contents of the exocrine glands of the ant subfamily Cerapachyinae, *Biochemical Systematics and Ecology*, 36:260-265.
165. Defensive chemicals of two species of *Trachypachus* MOTSCHULSKI, Attygalle A.B, Wu X, Ruzicka J, Rao S, Garcia S, Herath K, Meinwald J, Maddison D.R, and Will K.W, *Journal of Chemical Ecology*, 30(3): 577-587.
166. Eelen D, Borgesen L and Billen J, 2006. Functional morphology of the postpharyngeal gland of queens and workers of the ant *Monomorium pharaonis* (L) , *Acta Zoologica* (Stockholm) 87: 101-111.
167. Ralston J.P, Rapini R.P, Josey MD, 2010. Fire Ant Bites, *Emergency medicine*.
168. Jones TH, Roberston HG, Snelling RR, 2003. The venom alkaloids from some African *Monomorium* species, *J Chem Ecol*, 29 (12): 2721-2727.

169. Lemaire M, Lange C, Bazire M, Cassier P, Clement J L, Escoubas P, Basselier J J, 1988. Alkaloid venom of European ants in the genus *Monomorium*. Site of synthesis, identification and quantification, *Exp Biol*, 48 (1): 27-40.
170. Morgan E D, Jungnickel H, Keegans S J, Do Nascimento RR, Billen J, Gobin B, Ito F, 2003. Comparative survey of abdominal gland secretions of the ant subfamily Ponerinae, *J Chem Ecol*, 29(1): 95-114.
171. Billen J, 2009. Occurrence and Structural organization of the exocrine glands in the legs of Ants, *Arthropod Struct Dev*. 38(1): 2-15.
172. Moreau C.S, 2014. A practical guide to DNA extraction, PCR, and gene-based DNA sequencing in Insects, *HALTERES*, 5: 32-42.
173. Blaimer B. B, 2012. A subgeneric revision of *Crematogaster* and discussion of regional species-groups (Hymenoptera: Formicidae), *Zootaxa*, 3482: 47-67.
174. Souza E, Follett P.A, Price D.K, and Stacy E.A, 2008. Field Suppression of the Invasive Ant *Wasmannia auropunctata* (Hymenoptera: Formicidae) in a Tropical fruit Orchard in Hawaii, *J. econ. Entomol.* 101(4) : 1068-1074.
175. Vanderwoude C and Nadeau B, 2009. Application methods for paste bait formulations in control of Ants in arboreal situations, *PROC. HAWAIIAN ENTOMOL. SOC.* 41: 113-119.
176. Watanasit S and Jantarit S, 2006. The ant nest of *Crematogaster rogenhoferi* (Mayr, 1879) at Tarutao national park, Satun Province, Southern Thailand, *J.Sci. Technol.*, 2894: 723-730.
177. BANFER G, MOOG U, FIALA B, MOHAMED M, WEISING K and BLATTNERS F.R, 2006. A chloroplast genealogy of myrmecophytic *Macaranga* species (Euphorbiaceae) in Southeast Asia reveals hybridization, variance and long-distance dispersals, *Molecular Ecology*, 15(14): 4440-4454.
178. Kang J.D, Kim S.J, Youn N.H, Kim B.J, Park S.D, 2004. A case of Ant Sting by *Crematogaster* *Ann Dermatol* 16(2), 87-89.
179. Hosoishi S, 2015. Revision of the *Crematogaster ranavalonae* – group in Asia, with description of two new species (Hymenoptera, Formicidae), *Journal of Hymenoptera Research*, 42: 63-92.
180. Yamane S and Terayama M, 1999. A new species of the genus *Pristomyrmex* MAYR from Japan, and a proposal of a New synonym of species in the genus *Camponotus* MAYR (Hymenoptera; Formicidae), *Mem. Myrmecol. Soc. Jpn.*, 1 : 17-24.
181. Ng'endo R.N. <sup>1a\*</sup>, Osiemo Z.B. <sup>2b</sup>, Brandl R<sup>c</sup>, 2013. DNA barcodes for species identification in the hyperdiverse ant genus *Pheidole* (Formicidae: Myrmicinae), *Journal of Insect Science*, 13(27).
182. Lapolla<sup>1</sup> J. S., Brady S. G. <sup>2</sup> and Shattuck S. O. <sup>3</sup>, 2010. Phylogeny and Taxonomy of the *Prenolepis* genus-group of ants (Hymenoptera: Formicidae), *Systematic Entomology*, 35:118-131.
183. Mele P.V, 2008. A historical review of research on the weaver ant *Oecophylla* in biological control, *Agricultural and Forest Entomology*, 10: 13-22.
184. Dwomoh<sup>1\*</sup> E.A, Ackonor<sup>1</sup> J.B and Afun<sup>2</sup> J.V.K, 2008. Survey of insect species associated with Cashew (*Anacardium occidentale* Linn.) and their distribution in Ghana, *African journal of Agricultural Research* 3(3):205-214.
185. Crozier R.H, Newey P.S, Schluns E.A and Robson S.K.A, 2010. A masterpiece of evolution- *Oecophylla* weaver ants (Hymenoptera: Formicidae), *Myrmecological News*, 13: 57-71.

186. Abdulla N.R, Rwegasira G.M, Jensen K.M.V, Mwatawala M.W, Offenberg J, 2016. Potential of *Oecophylla longinoda* Latreille (Hymenoptera: formicidae) in managing major insect pests in organic cashew production systems, *Org. Agr.* 1-10.
187. Offenberg J, 2014. The use of artificial nests by weaver ants : A preliminary field observation, 6: 119-128.
188. Anato F.M<sup>\*1</sup>, Wargui R.B<sup>\*1</sup>, Sinzogan A.A.C, Offenberg J, Adandonon A, Vayssieres J.F and Kossou D.K, 2015. Reducing losses inflicted by insect pests on cashew, using weaver ants as a biological control agent, 17:285-291.
189. Tsuji K, Hasyim A, Harlion and Nakamura K, 2004. Asian weaver ants, *Oecophylla smaragdina*, and their repelling of pollinators, 19:669-673.
190. Bhattacharya A, 2004. Flower visitors and fruitset of *Anacardium occidentale* , *Ann.Bot.Fennici*, 41: 385-392.
191. Peng R, Lan L.P, and Christian K, 2014. Weaver Ant role in Cashew Orchards in Vietnam, *Biological and Microbial control*, 107(4): 1331-1338.
192. Abdulla N.R, Rwegasira G.M, Jensen K.M.V, Mwatawala M.W, Offenberg J, 2016. Potential of *Oecophylla longinoda* Latreille (Hymenoptera: Formicidae) in managing major insect pests in organic Cashew production systems, *Org. Agr.*
193. Crozier R.H, Newey P.S, Schuluns E.A, and Robson S.K.A, 2010. A masterpiece of evolution- *Oecophylla* weaver ants (Hymenoptera: Formicidae), *Myrmecological News*, 13: 57-71.
194. Dwomoh E.A, Ackonor J.B and Afun J.V.K , 2008. Survey of insect species associated with Cashew (*Anacardium occidentale* Linn.) and their distribution in Ghana, *African journal of Agricultural Research*, 3(3): 205-214.
195. Mele P.V, 2008. A historical review of research on the weaver ant *Oecophylla* in biological control, 10: 13-22.
196. Vanitha K, Bhat P.S, Raviprasad T.N, and Srikumar K.K, 2015. Species composition of Ants in Cashew plantations and their interrelationships with Cashew, *Proceedings of the national academy of sciences , India section B: Biological sciences*, 1-11.
197. Rickson F.R and Rickson M.M, 1998. The Cashew nut, *Anacardium occidentale* (Anacardiaceae), and its perennial association with Ants: extrafloral nectar location and the potential for Ant defense, *American journal of Botany*, 85(6): 835-849.
198. Freitas B.M and Paxton R.J, 1996.The role of wind and insects in Cashew (*Anacardium occidentale*) pollination in NE Brazil, *Journal of Agricultural Science, Cambridge*, 126: 319-326.
199. Bharti H and Silla S, 2011. Notes on life history of *Oecophylla smaragdina* (Fabricus) and its potential as biological control agent. *Halteres*, 3: 58-64.
200. Peng R K, Christian K and Gibb K, 2015. Distribution of the Green ant, *Oecophylla smaragdina* (F) (Hymenoptera: Formicidae), in relation to native vegetation and the insect pests in Cashew plantations in Australia, 43(3): 203-211.
201. Peng R K, Christian K, 2004. The weaver ant, *Oecophylla smaragdina* (Hymenoptera: Formicidae), an effective biological control agent of the red-banded thrips, *Selenothrips rubrocinctus* (Thysanoptera: Thripidae) in Mango crops in the Northern Territory of Australia, *International journal of pest management*,50(2): 107-114.

202. Peng R K, Christian K and Gibb K, 2005. Ecology of the fruit spotting bug, *Amblyopelta lutescens lutescens* Distant (Hemiptera: Coreidae) in Cahew plantations, with particular references to the potential for its biological control, Australian journal of Entomology, 44: 45-51.
203. Ativor I.N, Nuamah K A, Billah M K, and Ofori D O, 2012. Weaver Ant, *Oecophylla longinoda* (Latreille) (Hymenoptera: Formicidae) activity reduces fruit fly damage in Citrus orchards, Journal of agricultural Science and technology, A(20: 449-458.
204. Itterebeeck J V, Sivongxay N, Praxaysombath B, and Huis A V, 2015. Preliminary observations on gravid queen protection in *Oecophylla smaragdina* : Evacuation and retinue function, 7: 155-158.
205. Chapuisat M and Keller I, 2002. Division of labour influences the rate of ageing in weaver ant workers, The royal society, 269: 909-913.
206. Wargui R, Offenber J, Sinzogan A, Adandonon A, Kossou D and Vayssieres J F, 2015. Comparing different methods to assess weaver ant abundance in plantation trees, Asian Myrmecology, 7: 159-170.
207. Basnagala S, Wijesekara G.A.W and Wijayagunasekara H N P, 2002. Review of the subfamily Bryocorinae (Heteroptera: Miradae) of Sri Lanka, 14: 154-164.
208. Verghese A, Jayanthi P D K, Sreedevi K, Devi K S and Pinto V, 2013. A quick and non-destructive population estimate for the weaver ant *Oecophylla smaragdina* Fab. (Hymenoptera : Formicidae), Current Science, 104(5): 641-645.
209. Wijetunge P.M.A.P.K , Ranaweera B, 2015. Potential of Weaver Ant (*Oecophylla smaragdina* F.) for the Management of *Heloprltis antonii* Sign. In Cashew (*Anacardium occidentale* L.), Proceedings of International symposium on Cashew.
210. Rickson F R and Rickson M M, 1998. The Cashew nut , *Anacardium occidentale* (Anacardiaceae), and its perennial association with Ants: extrafloral nectar location and the potential for Ant defense. American Journal of Botany, 85(6): 835-849.
211. Offenber J, Peng R and Nielsen M G, 2012. Development rate and brood production in haplo- and pleometrotic colonies of *Oecophylla smaragdina* , Insectes Sociaux, 59: 307-311.
212. Peeters C and Andersen A N, 1998. Cooperation between dealate queens during colony foundation in the green tree Ant, *Oecophylla smaragdina*, 96: 40-44.